





Healthification

Percent Of Obese (BMI > 30) In U.S. Adults



State	%	State	%	State	%	State	%
Alabama	31.0	Illinois	26.5	Montana	23.2	Rhode Island	24.6
Alaska	24.8	Indiana	29.5	Nebraska	27.2	South Carolina	29.4
Arizona	25.5	Iowa	27.9	Nevada	25.8	South Dakota	29.6
Arkansas	30.5	Kansas	28.1	New Hampshire	25.7	Tennessee	32.3
California	24.8	Kentucky	31.5	New Jersey	23.3	Texas	28.7
Colorado	18.6	Louisiana	33.0	New Mexico	25.1	Utah	23.5
Connecticut	20.6	Maine	25.8	New York	24.2	Vermont	22.8
Delaware	27.0	Maryland	26.2	North Carolina	29.3	Virginia	25.0
Washington DC	19.7	Massachusetts	21.4	North Dakota	27.9	Washington	26.4
Florida	25.2	Michigan	29.6	Ohio	28.8	West Virginia	31.1
Georgia	27.2	Minnesota	24.6	Oklahoma	31.4	Wisconsin	28.7
Hawaii	22.3	Mississippi	34.4	Oregon	23.0	Wyoming	24.6
Idaho	24.5	Missouri	30.0	Pennsylvania	27.4		

The data shown in this map is the most recent collected through the CDC's Behavioral Risk Factor Surveillance System (BRFSS). Each year, state health departments use standard procedures to collect data through a series of monthly telephone interviews with U.S. adults. Prevalence estimates generated for the maps may vary slightly from those generated for the states by the BRFSS as slightly different analytic methods are used.

